

FORM PTO-1390 (Modified)
(REV 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTORNEY'S DOCKET NUMBER

TRANSMITTAL LETTER TO THE UNITED STATES

MIT-C205

DESIGNATED/ELECTED OFFICE (DO/EO/US)

U.S. APPLICATION NO. (IF KNOWN, SEE 37 CFR

CONCERNING A FILING UNDER 35 U.S.C. 371

10/070161

INTERNATIONAL APPLICATION NO.

INTERNATIONAL FILING DATE

PRIORITY DATE CLAIMED

PCT/JP00/06187

11 SEPT 2000

13 SEPT 1999

TITLE OF INVENTION

KITS AND METHODS FOR DETECTING OR MEASURING ANALYTES

APPLICANT(S) FOR DO/EO/US

Yuichi OKU, Hisanori KAMIYA, Kumiko SHINOHARA, Yusuke SIBAHARA and Yoshihiko UESAKA

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (24) indicated below.
4. ☐ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371 (c) (2))
 - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
 - b. ☒ has been communicated by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
- ☒ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
 - a. ☒ is attached hereto.
 - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
- ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371 (c)(3))
 - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
 - b. ☐ have been communicated by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
- ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
- ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371 (c)(4)).
- ☐ An English language translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371 (c)(5)).
11. ☐ A copy of the International Preliminary Examination Report (PCT/IPEA/409).
12. ☐ A copy of the International Search Report (PCT/ISA/210).

Items 13 to 20 below concern document(s) or information included:

13. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
14. ☒ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
15. ☒ A **FIRST** preliminary amendment.
16. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
17. ☐ A substitute specification.
18. ☐ A change of power of attorney and/or address letter.
19. ☒ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
20. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
21. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
22. ☐ Certificate of Mailing by Express Mail
23. ☒ Other items or information:

Four (4) sheets of formal drawings
Form PCT/IB/308

Statement Pursuant to 37 CFR 1.821(f)

APPLICATION NO. (IF KNOWN, SEE 37 CFR

10/070161

INTERNATIONAL APPLICATION NO.

PCT/JP00/06187

ATTORNEY'S DOCKET NUMBER

MIT-C205

24. The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)) :

- ☐ Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO \$1040.00
- ☒ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO \$890.00
- ☐ International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO \$740.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) \$710.00
- ☐ International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) \$100.00

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$890.00

Surcharge of \$130.00 for furnishing the oath or declaration later than months from the earliest claimed priority date (37 CFR 1.492 (e)).

☐ 20 ☐ 30

\$0.00

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	25 - 20 =	5	x \$18.00
Independent claims	6 - 3 =	3	x \$84.00

\$90.00

\$252.00

Multiple Dependent Claims (check if applicable).

☐

\$0.00

TOTAL OF ABOVE CALCULATIONS =

\$1,232.00

☐ Applicant claims small entity status. See 37 CFR 1.27). The fees indicated above are reduced by 1/2.

\$0.00

SUBTOTAL =

\$1,232.00

Processing fee of \$130.00 for furnishing the English translation later than months from the earliest claimed priority date (37 CFR 1.492 (f)).

☐ 20 ☐ 30

+

\$0.00

TOTAL NATIONAL FEE =

\$1,232.00

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31) (check if applicable).

☒

\$40.00

TOTAL FEES ENCLOSED =

\$1,272.00

Amount to be:

refunded

\$

charged

\$

- a. ☐ A check in the amount of _____ to cover the above fees is enclosed.
- b. ☐ Please charge my Deposit Account No. _____ in the amount of _____ to cover the above fees. A duplicate copy of this sheet is enclosed.
- c. ☐ The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. _____. A duplicate copy of this sheet is enclosed.
- d. ☒ Fees are to be charged to a credit card. **WARNING:** Information on this form may become public. **Credit card information should not be included on this form.** Provide credit card information and authorization on PTO-2038.

NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.

SEND ALL CORRESPONDENCE TO:

George A. Loud, Esq.
LORUSSO & LOUD
3137 Mount Vernon Avenue
Alexandria, VA 22305
(703) 739-9393



30132

PATENT TRADEMARK OFFICE

SIGNATURE

George A. Loud

NAME

25,814

REGISTRATION NUMBER

March 4, 2002

DATE

10/070161

JC19 Rec'd PCT/PTO 04 MAR 2002

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:

OKU et al

Serial No.:

Filed: March 4, 2002

For: KITS AND METHODS FOR DETECTING AND
MEASURING ANALYTES


STATEMENT PURSUANT TO 37 CFR 1.821(f)

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

The undersigned declares that the paper copy of the "Sequence Listing" submitted
herewith has the same content as the computer readable copy (diskette) submitted herewith.

Respectfully submitted,



George A. Loud
Reg. No. 25,814

Dated: March 4, 2002

LORUSSO & LOUD
3137 Mount Vernon Avenue
Alexandria, VA 22305

(703) 739-9393

20070161-000401

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:

OKU et al

Serial No.:

Filed: March 4, 2002

For: KITS AND METHODS FOR DETECTING AND
MEASURING ANALYTES

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Please amend the captioned application as follows:

IN THE CLAIMS:

Please rewrite claims 4, 6, 8-17 and 20-22 as follows:

4. (Amended) The kit according to claim 1, wherein the substance R1 has a plurality of binding points with respect to the substance B1, and a plurality of the ligands L1 are bound to the substance R1 by the medium of the substance B1.

6. (Amended) The kit according to claim 1, wherein the substance R2 has a plurality of binding points with respect to the substance B2, and a plurality of the ligands L2 or ligands L3 are bound

to the substance R2 by the medium of the substance B2.

8. (Amended) The kit according to claim 1, wherein the solid phase conjugate is at least independent of the receptor II.

9. (Amended) The kit according to claim 1, wherein the analyte A having bivalent or higher binding capability is a substance selected from the group consisting of DNAs, RNAs, antigens, and antibodies.

10. (Amended) The kit according to claim 1, wherein the ligand L1, ligand L2 or ligand L3 is a substance selected from the group consisting of DNA, RNA, antigen, antibody, lectin, glycoprotein, and sugar.

11. (Amended) The kit according to claim 1, wherein the ligand L1 and the ligand L2 are the same substance.

12. (Amended) The kit according to claim 1, wherein the ligands L1 and L2 and the ligand L3 are substances having different sequences with one another.

13. (Amended) The kit according to claim 1, wherein the binding capability between the substance B1 and the substance R1 or between the substance B2 and the substance R2 is represented by a dissociation constant of from 10^{-8} to 10^{-16} (M).

14. (Amended) The kit according to claim 1, wherein the substance B1 and/or the substance B2 is biotin.

15. (Amended) The kit according to claim 1, wherein the substance B1 and/or the substance B2 is a substance selected from the group consisting of DNA, RNA, antigen, antibody, lectin, glycoprotein, and sugar.

16. (Amended) The kit according to claim 1, wherein the substance R1 and/or the substance R2 is a substance selected from the group consisting of streptavidin and avidin.


17. (Amended) The kit according to claim 1, wherein the substance R1 and/or the substance R2 is a substance selected from the group consisting of antigen, antibody, DNA, RNA, lectin, glycoprotein, and sugar.

20. (Amended) The kit according to claim 1, wherein the bond element B3 is bondable to the anti-bond element R3 by complementary binding of at least part of DNAs or RNAs.

21. (Amended) The kit according to claim 1, wherein the marker M is a substance selected from the group consisting of coloring dye, fluorescent dye, luminescent substance, metal colloid, latex, liposome, radioactive isotope, enzyme, DNA, and RNA.

22. (Amended) The kit according to claim 1, wherein the solid phase is a substance selected from the group consisting of polystyrene, nitrocellulose, nylon, cellulose, and glass.

Respectfully submitted,



George A. Loud
Reg. No. 25,814

Dated: March 4, 2002

LORUSSO & LOUD
3137 Mount Vernon Avenue
Alexandria, VA 22305

(703) 739-9393

4. (Amended) The kit according to claim 1 [or 3], wherein the substance R1 has a plurality of binding points with respect to the substance B1, and a plurality of the ligands L1 are bound to the substance R1 by the medium of the substance B1.
6. (Amended) The kit according to claim 1 [or 2], wherein the substance R2 has a plurality of binding points with respect to the substance B2, and a plurality of the ligands L2 or ligands L3 are bound to the substance R2 by the medium of the substance B2.
8. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the solid phase conjugate is at least independent of the receptor II.
9. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the analyte A having bivalent or higher binding capability is a substance selected from the group consisting of DNAs, RNAs, antigens, and antibodies.
10. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the ligand L1, ligand L2 or ligand L3 is a substance selected from the group consisting of DNA, RNA, antigen, antibody, lectin, glycoprotein, and sugar.
11. (Amended) The kit according to claim 1 [or 3], wherein the ligand L1 and the ligand L2 are the same substance.

12. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the ligands L1 and L2 and the ligand L3 are substances having different sequences with one another.

13. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the binding capability between the substance B1 and the substance R1 or between the substance B2 and the substance R2 is represented by a dissociation constant of from 10^{-8} to 10^{-16} (M).

14. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the substance B1 and/or the substance B2 is biotin.

15. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the substance B1 and/or the substance B2 is a substance selected from the group consisting of DNA, RNA, antigen, antibody, lectin, glycoprotein, and sugar.

16. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the substance R1 and/or the substance R2 is a substance selected from the group consisting of streptavidin and avidin.

17. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the substance R1 and/or the substance R2 is a substance selected from the group consisting of antigen, antibody, DNA, RNA, lectin, glycoprotein, and sugar.

20. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the bond element B3 is bondable to the anti-bond element R3 by complementary binding of at least part of DNAs or RNAs.

21. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the marker M is a substance selected from the group consisting of coloring dye, fluorescent dye, luminescent substance, metal colloid, latex, liposome, radioactive isotope, enzyme, DNA, and RNA.

22. (Amended) The kit according to claim 1 [any one of claims 1 to 3], wherein the solid phase is a substance selected from the group consisting of polystyrene, nitrocellulose, nylon, cellulose, and glass.

DESCRIPTION

KITS AND METHODS FOR DETECTING OR MEASURING ANALYTES

5 1. Technical Field

10 The present invention relates to kits and methods for detecting or measuring an analyte having bivalent or higher binding capability and contained in a liquid sample. The kits each comprise a ligand-marker complex, a ligand-bond element complex, and a solid phase. The ligand-marker complex comprises a ligand which specifically binds to the analyte and is labeled with a marker directly or indirectly. The ligand-bond element complex comprises a ligand and a bond element, which ligand specifically binds to the analyte and which bond element serves to bind the ligand to the solid phase. The solid phase comprises an anti-bond element which allows the solid phase to capture the ligand-bond element complex.

15 2. Background Art

20 Known methods for detecting or measuring an antibody or another substance in a liquid sample, which substance has bivalent or higher binding capability to a ligand, include the following method. In the method, a ligand labeled with a marker such as an enzyme, and a solid phase bound to a ligand are previously prepared, and the ligand labeled with the marker is allowed to react with an antibody in a liquid sample. The resulting solution mixture is then allowed to react with the ligand bound to the solid phase to thereby determine the

25

presence or absence of the marker bound to the solid phase or to measure the amount thereof (e.g., Japanese Patent Application Publication No. 9-229938). According to this method, the antibody reactive with respect to the identical
5 ligand previously reacts with the labeled ligand and becomes unreactive with the solid phased ligand, and the resulting sensitivity is insufficient in many cases.

For this reason, for example, Japanese Patent No. 2532788 discloses a method for determining the presence or the amount of
10 an immunoreactive analyte (G). In this method, a capture moiety (R) is bound to a solid phase, a first batch (M-I1) of a first particle is bound to a second batch (I2-B) of a second particle through the medium of the immunoreactive analyte (G) and thereby yields a complex (M-I1-G-I2-B). The first batch (M-
15 I1) comprises a first immunological substance (I1) bound to a detectable isotope (M) and is dispersed. The second batch (I2-B) comprises a second binding product of a second immunoreactive substance (I2) and a species (B) capable of being captured and is dispersed. The formed complex (M-I1-G-
20 I2-B) is allowed to react with the capture moiety (R) bound to the solid phase to thereby determine the presence or amount of the immunoreactive analyte. According to this method, when the first immunological substance (I1) is identical to the second immunological substance, the first immunological substance (I1)
25 and the detectable isotope (M) are not bound to the immunoreactive analyte (G), but the first immunological substance (I1) and the second immunological substance (I2)

concurrently react with the immunoreactive analyte (G). Thus, the method exhibits a high sensitivity.

Japanese Patent Application Publication No. 10-253632 discloses a method for evaluating an analyte based on the presence or absence or the amount of a marker (M). In this method, an oligonucleotide is solid-phased, a ligand (L) labeled with an oligonucleotide (ON') being complementary to the oligonucleotide (ON) (the resulting oligonucleotide-labeled ligand is ON'-L, e.g., ON'-antigen or ON'-antibody) and a marker-labeled ligand (M-L) are allowed to react with the analyte (A) being capable of binding to the ligand (L) and thereby yield a complex (ON'-L-A-L-M). The complex is then bound to the solid phase by complementary binding between oligonucleotides contained in the oligonucleotide-binding solid phase and the complex, respectively, to determine the presence or absence or the amount of the marker (M). This method exhibits a high sensitivity as in the method described in Japanese Patent No. 2532788, since the analyte (A) and the marker-labeled ligand (M-L) do not previously form a complex, but ON'-L and M-L concurrently react with the analyte (A). According to the method in question, plural analytes can concurrently be measured by changing the sequences of the oligonucleotide to be solid-phased and of the oligonucleotide to be bound to the corresponding ligand (L).

To detect or measure an analyte with a high sensitivity as in the methods described in Japanese Patent No. 2532788 and Japanese Patent Application Publication No. 10-253632, a first

or second immunoreactive substance or a ligand must be labeled with a bond element of some kind.

However, when the immunoreactive substance or the ligand (L) is directly labeled with the bond element, the ligand (L) changes in physical properties and tends to become insoluble in a purification step of the labeled ligand (L) during labeling process. Accordingly, the recovery of the resulting bond element-labeled ligand (e.g., an antigen labeled with the bond element) significantly decreases in some cases.

When a low molecular weight peptide as a ligand (L) is directly labeled, the labeled peptide can be recovered but may show decreased immunochemical reactivity. This is probably caused by steric hindrance. In addition, when the low molecular weight peptide is to be labeled with a metal colloid or a colored latex, it may not be labeled because of its low molecular weight.

Japanese Patent No. 2501960 discloses a reagent. The reagent comprises a substance P2 (e.g., avidin or streptavidin) that is labeled with a marker and has specific binding capability, a substance P2 that is not labeled with a marker and has specific binding capability, a compound P1-R1, and a solid phase bound to the substance P2 that is not labeled with a marker. The compound P1-R1 comprises a component P1 (e.g., biotin) having monovalent binding capability to P2, and a ligand R1 bound to P1. The reagent cannot significantly be controlled in sensitivity, since the amount of the ligand per molecule of the complex cannot be controlled, or the reaction with the solid

phase yields a strong bond derived from the bond between biotin and avidin. The reagent may also react with a trace quantity of an immunity substance originated from another factor and may yield unsuitable conclusions.

5 Accordingly, an object of the present invention is to provide a kit and a method for detecting or measuring an antibody or another analyte contained in a liquid sample, which analyte has bivalent or higher binding capability to a ligand. The kit can easily detect or measure the analyte, can be
10 prepared in a high yield and can easily label a target substance regardless of whether it is a low molecular weight ligand, a high molecular weight ligand or a ligand constituting part of a protein. The kit can produce a ligand-marker complex in a high
15 yield even when, for example, a ligand of a low molecular weight compound is labeled with a marker. In addition, the amount of the ligand in the kit can be controlled, or the binding capability between the solid phase and the complex can be changed, and therefore the sensitivity of a constitutional reagent can be controlled.

20

DISCLOSURE OF INVENTION

The present invention provides the following kits to solve the above problems.

25 A kit of a first type according to the present invention is a kit for detecting or measuring an analyte having bivalent or higher binding capability. The kit comprises the following receptor I, receptor II, and solid phase conjugate for

capturing a complex comprising the receptor I, the receptor II and the analyte:

1) a receptor I represented by a compound L1-B1-R1-M comprising a compound R1-M bound to a compound L1-B1, the compound R1-M comprising a substance R1 bound to a marker M, the substance R1 being capable of binding to a substance B1, the compound L1-B1 comprising a ligand L1 bound to the substance B1, the ligand L1 being capable of binding to the analyte A;

2) a receptor II represented by a compound L2-B2-R2-B3 comprising a compound L2-B2 previously bound to a compound R2-B3, the compound L2-B2 comprising a ligand L2 and a substance B2 introduced into the ligand L2, the ligand L2 being capable of binding to the analyte A, the substance B2 having different binding capability from analyte A, the compound R2-B3 comprising a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3 having binding capability different from the substance B2; and

3) solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3.

A kit of a second type according to the present invention is a kit for detecting or measuring an analyte A having bivalent or higher binding capability to a ligand L3 constituting part of a protein P. The kit comprises the following receptor I, receptor II, and solid phase conjugate for capturing a complex

comprising the receptor I, the receptor II and the analyte:

1) a receptor I represented by a compound P-M comprising the protein P bound to a marker M;

2) a receptor II represented by a compound P-B2-R2-B3 or L3-B2-R2-B3 comprising a compound P-B2 or L3-B2 bound to a compound R2-B3, the compound P-B2 or L3-B2 comprising the protein P or the ligand L3 and a substance B2 introduced into the protein P or the ligand L3, the substance B2 having different binding capability from analyte A, the compound R2-B3 comprising a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3 having binding capability different from the substance B2; and

3) solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3.

In the kits of the present invention, the solid phase conjugate is at least independent of the receptor II. Specifically, as assembly in the kits of the present invention before measurement, the solid phase conjugate is independent of the receptor II and the receptor I, or the solid phase conjugate is contained in the same system with the receptor I but is independent of the receptor II.

A kit of a third type according to the present invention is a kit for detecting or measuring an analyte A having bivalent or higher binding capability, the kit comprising the following receptor I, receptor II, and solid phase conjugate for

capturing a complex comprising the receptor I, receptor II and analyte:

5 1) a receptor I represented by a compound L1-B1-R1-M comprising a compound R1-M bound to a compound L1-B1, the compound R1-M comprising a substance R1 bound to a marker M, the substance R1 being capable of binding to a substance B1, the compound L1-B1 comprising a ligand L1 bound to the substance B1, the ligand L1 being capable of binding to the analyte A;

10 2) a receptor II represented by a compound L2-B3, the compound L2-B3 comprising a ligand L2 previously bound to a bond element B3, the ligand L2 being capable of binding to the analyte A, the bond element B3 having different binding capability from analyte A; and

15 3) solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3.

20 The analyte A having bivalent or higher binding capability in the kits of the present invention can be selected from the group consisting of DNA, RNA, antigen, and antibody.

25 In the kits of the present invention, the ligand L1 and the ligand L2 may be the same substance (e.g., both are avidin or both are streptavidin) or may be different substances having different sequences (e.g., both are DNAs or RNAs having different sequences or are avidin and streptavidin).

The ligands L (L1, L2, L3) in the kits of the present invention can be selected from the group consisting of DNA, RNA,

antigen, antibody, lectin, glycoprotein, and sugar.

The substance R1 and/or the substance R2 in the kits of the present invention can be selected from the group consisting of streptavidin, avidin, antigen, antibody, DNA, RNA, lectin, glycoprotein, and sugar.

The substance R1 and the substance R2 in the kits of the present invention may be the same substance (e.g., both are avidin or both are streptavidin) or different substances (e.g., avidin and streptavidin).

The substance B1 and/or the substance B2 in the kits of the present invention can be a substance selected from the group consisting of DNA, RNA, antigen, antibody, lectin, glycoprotein, and sugar. Biotin is typically used as the substance B1 and/or the substance B2 in an embodiment of the kits of the present invention. In this case, the substance R1 and the substance R2 are both avidin, both streptavidin, or a combination of avidin and streptavidin.

In the kits of the present invention, the binding capability between the substance B1 and the substance R1 or between the substance B2 and the substance R2 is preferably represented by a dissociation constant of from 10^{-8} to 10^{-16} (M).

The marker in the kits of the present invention may be a substance selected from the group consisting of coloring dye, fluorescent dye, luminescent substance, metal colloid, latex, liposome, radioactive isotope, enzyme, DNA, and RNA.

The solid phase for use in the present invention is preferably a substance selected from the group consisting of

polystyrene, nitrocellulose, nylon, cellulose, and glass.

A method for detecting or measuring an analyte A according to the present invention using the kit of the first type is a method for detecting or measuring an analyte A having bivalent
5 or higher binding capability. The method comprises the steps of:

bringing 1) the analyte A into contact with:

2) a receptor I represented by a compound L1-B1-R1-M comprising a compound R1-M bound to a compound L1-B1, the
10 compound R1-M comprising a substance R1 bound to a marker M, the substance R1 being capable of binding to a substance B1, the compound L1-B1 comprising a ligand L1 bound to the substance B1, the ligand L1 being capable of binding to the analyte A; and

3) a receptor II represented by a compound L2-B2-R2-B3 comprising a compound L2-B2 previously bound to a compound R2-B3,
15 B3, the compound L2-B2 comprising a ligand L2 and a substance B2 introduced into the ligand L2, the ligand L2 being capable of binding to the analyte A, the substance B2 having different binding capability from analyte A, the compound R2-B3 comprising
20 a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3 having binding capability different from the substance B2, to react with one another to thereby form a complex;

4) allowing a solid phase conjugate to capture the formed
25 complex, the solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3

being capable of binding to the bond element B3, and

5) detecting or measuring the marker M in the captured complex.

5 A method for detecting or measuring an analyte A of the present invention using the kit of the second type is a method for detecting or measuring an analyte A having bivalent or higher binding capability to a ligand L3 constituting part of a protein P. The method includes the steps of:

bringing 1) the analyte A into contact with:

10 2) a receptor I represented by a compound P-M comprising the protein P bound to a marker M; and

3) a receptor II represented by a compound P-B2-R2-B3 or L3-B2-R2-B3 comprising a compound P-B2 or L3-B2 bound to a compound R2-B3, the compound P-B2 or L3-B2 comprising the protein P or
15 the ligand L3 and a substance B2 introduced into the protein P or the ligand L3, the substance B2 having different binding capability from analyte A, the compound R2-B3 comprising a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3
20 having binding capability different from the substance B2, to react with one another to thereby form a complex,

4) allowing a solid phase conjugate to capture the formed complex, the solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond
25 element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3, and

5) detecting or measuring the marker M in the captured

complex.

A method for detecting or measuring an analyte A of the present invention using the kit of the third type is a method for detecting or measuring an analyte A having bivalent or
5 higher binding capability. The method includes the steps of:

bringing 1) the analyte A into contact with:

2) a receptor I represented by a compound L1-B1-R1-M comprising a compound R1-M bound to a compound L1-B1, the compound R1-M comprising a substance R1 bound to a marker M,
10 the substance R1 being capable of binding to a substance B1, the compound L1-B1 comprising a ligand L1 bound to the substance B1, the ligand L1 being capable of binding to the analyte A; and

3) a receptor II represented by a compound L2-B3, the compound L2-B3 comprising a ligand L2 previously bound to a bond
15 element B3, the ligand L2 being capable of binding to the analyte A, the bond element B3 having different binding capability from analyte A, to react with one another to thereby form a complex,

4) allowing a solid phase conjugate to capture the formed
20 complex, the solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3, and

5) detecting or measuring the marker M in the captured
25 complex.

In the three methods for detecting or measuring the analyte A, the complex comprising the elements 1), 2) and 3) may be

formed in one step by concurrently proceeding reactions or may sequentially be formed in plural steps by changing the order of bringing the elements into contact with one another.

Embodiments of the compound L1-B1-R1-M (receptor I) for use
5 in the kit of the present invention of the first type include a compound comprising streptavidin or avidin as the substance R1 which is bound to a metal colloid or enzyme as the marker M to form R1-M, and a biotin-bound ligand as the compound L1-B1 bound to R1-M by action of specific binding reaction between biotin
10 and avidin or streptavidin. This compound is a novel substance.

The kit of the first type of the present invention can be applied to the case where the ligand L1 is soluble in water and can advantageously be applied specifically to the case where the
15 ligand L1 is highly insoluble. When such a highly insoluble ligand L1 is directly labeled, the resulting complex may become insoluble owing to the insolubility of the ligand L1 and may cause a nonspecific reaction. However, by binding the ligand with the substance R1 or R2 having relatively high
20 hydrophilicity, such as avidin or streptavidin, the resulting complex represented by the compound L1-B1-R1-M (receptor I) or the compound L2-B2-R2-B3 (receptor II) can become hydrophilic as a whole and can prevent occurrence of a nonspecific reaction.

25 In this connection, when the biotin-avidin reaction is used for preparing individual complexes constituting the kit, the condition for the preparation of the complexes can be

maintained constant regardless of the type of the ligand L1. This configuration can save extra time and effort. According to the present invention, the complex L2-B2-R2-B3 (receptor II) is prepared in advance. The amount of the ligand per molecule
5 of the complex can therefore be controlled and the sensitivity of the reagent can thereby be controlled.

The kit of the second type of the present invention is suitable when the ligand L2 is not a low molecular weight substance but a water-soluble polymer such as a water-soluble
10 protein. When such a protein is used as the ligand L2, the ligand L2 can directly be labeled with the marker M.

The kit of the third type of the present invention is suitable for measuring an oligonucleotide in DNA or RNA as the analyte A. In this case, the ligand L1 is an oligonucleotide
15 at least having a complementary sequence with respect to a sequence of one end of the oligonucleotide analyte A, and the ligand L2 is an oligonucleotide at least having a complementary sequence with respect to a sequence of the other end of the analyte A. In addition, the sequence of an end of the ligand L2
20 opposite to that bound to the analyte A is at least complementary with respect to the sequence of the oligonucleotide of the DNA or RNA bound to the solid phase.

BRIEF DESCRIPTION OF THE DRAWINGS

25 Fig. 1 illustrates an embodiment of components of the kit of the first type of the present invention. During storage of the kit, a component in one frame is stored in configuration

separated from with components in the other frame.

In Fig. 2, the analyte A is captured by the solid phase using the kit of the first type of the present invention.

Fig. 3 is a flowchart for the preparation of reagent
5 components of the kit of the first type of the present invention. Specifically, it is a preparation flowchart for the receptor I represented by the compound L1-B1-R1-M and for the receptor II represented by the compound L2-B2-R2-B3.

Fig. 4 illustrates an embodiment of components of the kit of
10 the second type of the present invention. During storage of the kit, a component in one frame is stored in configuration separated from with components in the other frame.

In Fig. 5, the analyte A is captured by the solid phase using the kit of the second type of the present invention.

Fig. 6 illustrates an embodiment of components of the kit of
15 the third type of the present invention. During storage of the kit, a component in one frame is stored in configuration separated from with components in the other frame.

In Fig. 7, the analyte A is captured by the solid phase
20 using the kit of the third type of the present invention.

In Fig. 8, the kit of the present invention comprising the
ligand L1 and the ligand L2 having different sequences from
each other is allowed to react with the analyte A to thereby
allow the solid phase conjugate to capture a complex comprising
25 the receptor I, the receptor II and the analyte.

BEST MODE FOR CARRYING OUT THE INVENTION

The technical significance of the present invention will be illustrated with reference to several embodiments of the invention.

Avidin or streptavidin satisfactorily binds to biotin with
5 an association constant of 10^{-15} (M), 100 folds or more higher than those of general immunoreactions. It has been reported that this bond dissociates within 15 minutes under the condition of 6-8 M guanidine, pH 1.5 and 120° C (Green, N.: Purification of Avidin, in Methods in Enzymology, XVIII,
10 (McCormick, D., and Wright, L., eds.), Academic Press, NY, 414 (1970)) to thereby form a complex that is very stable at room temperature.

An example of the kit of the first type of the present invention comprising avidin or streptavidin as R1 or R2 will be
15 illustrated. In this kit, use can be made of a receptor I represented by the compound L1-B1-R1-M or a receptor II represented by the compound L2-B2-R2-B3. The compound L1-B1-R1-M comprises the ligand L1 bound to the marker M and the compound L2-B2-R2-B3 comprises the ligand L2 bound to an oligonucleotide
20 as the bond element B3, each by action of strong bond between avidin or streptavidin as R1 or R2 and biotin or another component as B1 or B2.

In the preparation of the receptor I, initially, biotin as the substance B1 is introduced into the ligand L1 to form the
25 compound L1-B1, and the marker M is bound to avidin or streptavidin as the substance R1 to form the compound R1-M. Avidin or streptavidin R1 and biotin B1 in these compounds are

then bound to each other and thereby very easily and stably yield the complex (receptor I) represented by the compound L1-B1-R1-M. The above-prepared complex is substantially a substance comprising the ligand L1 labeled with the marker M.

5 The compound L1-B1-R1-M (receptor I) is the complex prepared according to the above method and comprises the marker M bound to, for example, avidin or streptavidin (R1) and the biotinylated ligand (the compound L1-B1) bound to avidin or streptavidin (R1). Accordingly, even when the ligand L1 is a
10 low molecular weight peptide or another substance that is susceptible to steric hindrance, it becomes less subject to such hindrance, as the marker M is bound to the ligand L1 through the medium of the high molecular weight avidin or streptavidin (R1). In addition, even when the ligand L1 is a
15 substance that cannot directly be bound to the marker M, the method can reliably produce the complex of the ligand L1 and the marker M.

When the ligand L3 is a relatively stable substance that is resistant to steric hindrance, that is, when the ligand L3
20 constitutes part of a protein, the marker M can directly be bound to the ligand L3 or to the protein P containing the ligand L3, and the resulting compound P-M can be used as a ligand-marker complex (receptor I) (the kit of the second type of the present invention).

25 When the substance B1 is identical to the substance B2, and the ligand L1 is identical to the ligand L2, the kit of the first type of the present invention can easily be prepared

10070161.030402

according to the following method. To prepare the receptor II, i.e., the compound (L2-B2-R2-B3) comprising the ligand L2 compounded with the bond element B3, the substance R2 is previously bound to the bond element B3 such as an oligonucleotide ON and thereby yields the compound R2-ON. The compound L1-B1 prepared in the preparation of the receptor I (L1-B1-R1-M), wherein L1-B1 is the same compound with L2-B2, is then allowed to react with the compound R2-ON and thereby yields a complex (receptor II) represented by a compound L2-B2-R2-ON. This method can prepare the complex comprising the ligand L2 and the nucleotide ON (bond element B3).

This method can reliably prepare the complex of the ligand L2 and the nucleotide ON (bond element B3) through the medium of the substance R2 (e.g., a substance selected from the group consisting of streptavidin, avidin, antigen, antibody, DNA, RNA, lectin, glycoprotein, and sugar) and the substance B2 even when the ligand L2 is a low molecular weight peptide or another substance that is difficult to labeling.

To compose the kit of the first type of the present invention, the compound R1-M and the compound R2-B3 (e.g., R2-ON) are prepared in advance, and the compound L1-B1 is used in common in the case where the compound L1-B1 is identical to the compound L2-B2, and the complex of the ligand L1 and the marker M, i.e., the receptor I represented by L1-B1-R1-M, and the complex of the ligand L2 and the bond element B3 (e.g., the oligonucleotide ON), i.e., the receptor II represented by the compound L2-B2-R2-B3, are then prepared as shown in the

flowchart in Fig. 3. By this procedure, the kit of the first type of the present invention can very easily be prepared.

Avidin or streptavidin for use as the substance R1 or the substance R2 has four binding sites per molecule with respect to biotin as the substance B1 or the substance B2. Accordingly, the introduction ratio of the ligand L1 or the ligand L2 can be changed in the range from one to four per molecule of avidin or streptavidin, and the reactivity and sensitivity can therefore be controlled by controlling the introduction ratio.

When the ligand L1 or the ligand L2 is a highly hydrophobic substance, and the marker M is directly introduced into the ligand L1 or the bond element B3 such as the oligonucleotide ON is directly introduced into the ligand L2, the physical properties further change, the hydrophobicity is further enhanced and thereby causes nonspecific bonds in many cases. However, when the ligand-biotin is bound to the avidin-marker or the avidin-bond element as in the kit of the first type of the present invention and when the ligand-marker complex or the ligand-bond element complex is prepared, the hydrophobicity of the ligand can be mitigated by action of binding with avidin having high hydrophilicity, thus eliminating nonspecific reactions due to hydrophobicity. In addition, the reactivity can also be controlled in this procedure by changing the introduction ratio of avidin to the ligand-biotin.

The above explanation is mainly on the kit of the first type of the present invention but can also be applied to the kit of the second type of the present invention as in the kit of the

first type.

The invented kits and methods for detection or quantification use the bond element B3 bound to the anti-bond element R3 to allow the solid phase to capture the complex. By
5 employing combinations of substances having various binding capabilities, the intensity of the binding capability can be varied, and thereby the sensitivity of the reagent can be controlled. For example, the binding reaction between lectin and a sugar exhibits reactivity much lower than that of the
10 biotin-avidin binding reaction. Accordingly, when the lectin and sugar are used as B3 and R3, the resulting kit does not react unless an antigen or antibody in a much more amount is contained in the sample than that in a kit using the biotin-avidin reaction.

15 When the bond element B3 and the anti-bond element R3 are oligonucleotides of RNA and/or DNA at least having partly complementary sequences and are bound to each other by action of their complementarity, the intensity of binding capability can be controlled, and thereby the sensitivity of the reagent can be
20 controlled by changing the lengths and/or homology of the complementary sequences of the oligonucleotides.

The invented kits and methods for detection or quantification can simultaneously detect or measure plural analytes by preparing plural combinations of oligonucleotides to
25 be solid-phased and corresponding oligonucleotides to be bound to corresponding ligands and selecting ligands that can specifically be bound to the plural analytes corresponding to

the individual combinations to thereby constitute a kit.

With reference to Fig. 8, the kit of the present invention comprising a ligand L1 and a ligand L2 having different sequences from each other is used and is allowed to react with an analyte A, and thereby the complex comprising the receptor I, the receptor II and the analyte is captured by the solid phase conjugate. Specifically, the analyte A shown in Fig. 8 is an oligonucleotide of an RNA or DNA. One oligonucleotide and the other having complementary sequences to both ends of the aforementioned oligonucleotide are used as the ligands L1 and L2, respectively. The components of the kit constituting the embodiment shown in Fig. 8 are the same with those of the kit of the first type of the present invention.

EXAMPLE 1

Synthesis of Oligonucleotides

The following oligonucleotides each having an amino group at five prime end were synthesized by using a DNA synthesizer available from The Perkin-Elmer Corporation, respectively. Some of the oligonucleotides were synthesized by Sawady Technology Co., Ltd.

Amino group-GAA TTC CCG GGG ATC CGT CG (hereinafter referred to as "Pair 1+")

Amino group-CGA CGG ATC CCC GGG AAT TC (hereinafter referred to as "Pair 1-")

Amino group-AAC GGA ATC TAA TCA GGA GG (hereinafter referred to as "Pair 8+")

Amino group-CCT CCT GAT TAG ATT CCG TT (hereinafter referred to

as "Pair 8-")

Amino group-CCG ACT ACA GAA GAG GAG AA (hereinafter referred to as "Pair 7+")

EXAMPLE 2

5 Preparation of Oligonucleotide-labeled IgG for Detection of Anti-HCV Antibody and TP

Pair 8- was introduced into a rabbit IgG. Specifically, 2 ml of 0.1 M sodium borate buffer containing 25 mg of the rabbit IgG was allowed to react with a 50- to 100-fold excess by mole of N-succinimidyl S-acetylthioacetate (SATA) (available from
10 Pierce Chemical Company) with respect to IgG at 37 ° C for 1 hour. After the reaction, Tris-HCl buffer and hydroxylamine were added to the reaction mixture to a final concentration of 0.1 M, respectively, and the resulting mixture was allowed to
15 react at 37° C for 1 hour. The mixture was then applied to a column filled with Sephadex G-25 (available from Amersham Pharmacia Biotech Company (Amersham Biosciences)) and thereby yielded an SH-introduced rabbit IgG. Separately, to 0.1 M 3-(N-morpholino)propanesulfonic acid (MOPS) buffer pH 7.0
20 containing 1000 nmol of Pair 8- dissolved therein and 5 mM EDTA, a 50-fold by mole excess of EMCS (available from Dojindo) with respect to the oligonucleotide was added and was allowed to react at 37 ° C for 1 hour. After the reaction, the resulting mixture was subjected to precipitation with ethanol and rinsing
25 according to a conventional procedure and thereby yielded a maleimide-group-introduced oligonucleotide. The maleimide-group-introduced oligonucleotide was allowed to react with the

SH-group-introduced IgG at 37 ° C for 1 hour, was then applied to a Ultrogel AcA 34 column (available from Biosepra) and thereby yielded a "Pair 8"-labeled rabbit IgG.

EXAMPLE 3

5 Preparation of colloidal gold-labeled Streptavidin

10 A total of 5 ml of colloidal gold having particle diameter of 40 nm (absorbance at 530 nm) was added to and was allowed to react with 1 ml of 2 mM borate buffer pH 9.0 containing 150 ug of streptavidin (available from Vector Laboratories). The resulting mixture was blocked with bovine serum albumin with a final concentration of 1%, a colloidal gold-labeled streptavidin was then recovered as a precipitate by centrifugation, was subjected to centrifugal washing with 1% bovine serum albumin solution, the resulting precipitate was suspended in the
15 solution and thereby yielded the target colloidal gold-labeled streptavidin.

EXAMPLE 4

Preparation of Oligonucleotide-labeled Streptavidin

20 To 0.1 ml of 0.2 M borate buffer pH 8.0 containing 2 mg of streptavidin dissolved therein, a 100-fold by mole excess of SATA with respect to the streptavidin was added and was allowed to react at 37° C for 1 hour. Tris buffer and hydroxylamine were then added to the resulting mixture to a final concentration of 0.1 M, respectively, and the resulting mixture
25 was allowed to react at 37° C for 30 minutes. The reaction mixture was then applied to a Sephadex G-25 column and thereby yielded an SH-group-introduced streptavidin. Separately, to

0.1 M MOPS buffer pH 7.0 containing 216 nmol of Pair 8+ dissolved therein and 5 mM EDTA, a 100-fold by mole excess of EMCS with respect to the oligonucleotide was added and was allowed to react at 37° C for 1 hour. After the reaction, the resulting mixture was subjected to precipitation with ethanol and washing according to a conventional procedure and thereby yielded a maleimide-group-introduced oligonucleotide. The maleimide-group-introduced oligonucleotide was allowed to react with the SH-group-introduced streptavidin at 37 ° C for 1 hour, was then applied to a Ultrogel AcA 34 column and thereby yielded a "Pair 8+"-labeled streptavidin.

EXAMPLE 5

Preparation of HCV Core 3 Peptide-binding Colloidal gold-labeled Streptavidin

A total of 200 ml of the colloidal gold-labeled streptavidin prepared in Example 3 was added to 5 ul of a biotin-introduced hepatitis C virus (HCV) core 3 peptide (available from Innogenetics NV) and was allowed to react at 37 ° C for 1 hour. The resulting colloidal gold complex was recovered as a precipitate by centrifugation, was centrifugally washed with a 1% bovine serum albumin solution, the resulting precipitate was suspended in the solution, and thereby yielded HCV core 3 peptide-binding colloidal gold-labeled streptavidin.

EXAMPLE 6

Preparation of HCV Core 3 Peptide-binding Oligonucleotide-labeled Streptavidin

To 200 ug of the oligonucleotide-labeled streptavidin

prepared in Example 4, 56 ug of the biotin-introduced HCV core 3 peptide was added and was allowed to react at 37 ° C for 1 hour, the resulting mixture was then subjected to ultrafiltration using YM30 (available from Millipore Corporation) and thereby yielded a HCV core 3 peptide-binding oligonucleotide-labeled streptavidin.

EXAMPLE 7

Preparation of "Pair 8"-labeled IgG-binding Nitrocellulose Membrane

10 The "Pair 8"-labeled rabbit IgG prepared in Example 2 was diluted to 1 mg/ml, was charged into XY3000 (the trade name of an XY dispensing system for the preparation of test strips, available from BioDot, Inc.) and was applied to a nitrocellulose membrane (available from Advanced Microdevices Pvt. Ltd. (mdi))
15 lined with poly ethylene terephthalate (PET). After drying, the resulting article was blocked with Blockace (the trade name of a blocking agent available from Snow Brand Milk Products Co., Ltd.) at room temperature for 3 hours, was washed with distilled water and was then dried. A cellulose filter paper
20 (available from Whatman Inc. under the trade name of WF1.5 or from Advanced Microdevices Pvt. Ltd. (mdi)) was attached to one end of the dried article, and a conjugate release pad (a glass filter available from Advanced Microdevices Pvt. Ltd. (mdi)) to the other end. In this procedure, part of them were overlapped
25 with each other. The resulting article was cut to 5 mm in width and was housed in a housing.

EXAMPLE 8

Reactivity in Anti-HCV Antibody Positive Serum

10070151 030402
A total of 100 ul of an anti-HCV antibody positive serum was added to 11.5 ul of a solution containing 5 ul of the HCV core 3 peptide-binding colloidal gold-labeled streptavidin prepared in Example 5, 150 ng of the HCV core 3 peptide-binding oligonucleotide-labeled streptavidin prepared in Example 6, 4.4% Tween 80, and 44 mM EDTA, the resulting mixture was mixed and was immediately applied to a sample supply opening of the "Pair 8"-labeled IgG-binding nitrocellulose membrane prepared in Example 7. A quarter of an hour later, a reddish violet line of the colloidal gold was observed in a portion where the "Pair 8"-labeled IgG was applied. In contrast, no line was observed when a serum that turned out to contain no anti-HCV antibody was added as a control. These results revealed that the method of the present invention can determine the presence of a specific antibody to a ligand.

EXAMPLE 9

Preparation of colloidal gold-labeled TP Antigen

A total of 5 ml of colloidal gold having a particle diameter of 40 nm was mixed with and was allowed to react with 1 ml of 2 mM borate buffer pH 9.0 containing 100 ug of a genetically modified Treponema pallidum (TP) 17K antigen (available from Lee Laboratories). The resulting mixture was then blocked with bovine serum albumin with a final concentration of 1%, was subjected to centrifugation and thereby yielded a colloidal gold-labeled TP antigen as a precipitate. The recovered colloidal gold-labeled TP antigen was centrifugally washed with

a 1% bovine serum albumin solution, the resulting precipitate was suspended in the solution and thereby yielded a colloidal gold-labeled TP antigen.

EXAMPLE 10

5 Preparation of Oligonucleotide-labeled Avidin

To 1 ml of 0.1 M phosphate buffer pH 7.0 containing 10 mg of
avidin dissolved therein, a 100-fold by mole excess of EMCS
with respect to the avidin was added and was allowed to react
at 37° C for 1 hour. The resulting mixture was then applied to
10 a Sephadex G-25 column and thereby yielded a maleimide-group-
introduced avidin. Separately, to 0.1 M MOPS buffer pH 7.0
containing 375 nmol of Pair 8+ dissolved therein and 5 mM EDTA,
a 100-fold by mole excess of SATA with respect to the
oligonucleotide was added and was allowed to react at 37° C for
15 1 hour. After the reaction, Tris buffer and hydroxylamine were
added to the resulting mixture to a final concentration of 0.1
M, respectively, and were allowed to react at 37° C for 30
minutes. The reaction mixture was then subjected to
precipitation with ethanol and rinsing according to a
20 conventional procedure and thereby yielded an SH-group-
introduced oligonucleotide. The SH-group-introduced
oligonucleotide was allowed to react with the maleimide-group-
introduced avidin at 37 ° C for 1 hour, the resulting mixture
was applied to Ultrogel AcA44 and thereby yielded "Pair 8+"-
25 labeled avidin.

EXAMPLE 11

Preparation of Biotin-labeled TP 17K Antigen

To 0.5 ml of 0.1 M phosphate buffer pH 7.0 containing 500 ug of a TP 17K antigen (available from Innogenetics NV) and 5 mM EDTA, a 100-fold by mole excess of NHS-biotin (available from Pierce Chemical Company) with respect to the TP 17K antigen was added and was allowed to react at 37° C for 1 hour. After the reaction, the resulting mixture was applied to a Sephadex G-25 column and thereby yielded a biotin-labeled TP 17 antigen with a recovery of TP 17K antigen of 75%.

EXAMPLE 12

10 Preparation of Oligonucleotide-labeled TP 17K Antigen Through Reaction of Biotinylated TP 17K Antigen and Oligonucleotide-labeled Avidin

The oligonucleotide-labeled avidin prepared in Example 10 was mixed with the biotin-labeled TP 17K antigen prepared in Example 11 in a ratio of the former to the latter of 1:8, 1:4, 1:2, 1:1, and 1:0.5, respectively, and was allowed to react at 37° C for 30 minutes. The resulting mixtures were stored at 4 ° C.

In these mixtures, the recovery of the TP 17K antigen was 75% since they were prepared merely by mixing the oligonucleotide-labeled avidin prepared in Example 10 and the biotin-labeled TP 17K antigen prepared in Example 11.

EXAMPLE 13

25 Reactivity of Oligonucleotide-labeled TP 17K Antigen with Varying Different Mixing Ratios

The reactivity of each of the oligonucleotide-labeled TP 17K antigens with different mixing ratios prepared in Example 12

was evaluated. Specifically, 2 ul each of the oligonucleotide-labeled TP 17K antigens prepared in Example 12 was added to 5 ul of the colloidal gold-labeled TP antigen prepared in Example 9, 5 ul of 1 mg/ml biotin, and 90 ul of an anti-TP antibody negative serum, respectively. Two bottles of this reaction system were prepared, an anti-TP antibody positive serum was added only to one reaction system, and each of the two reaction systems was then rapidly mixed. The total amount of each of the resulting mixtures was applied to the sample supply opening of the housing including the "Pair 8-"-labeled IgG-binding nitrocellulose membrane prepared in Example 7.

Twenty minutes after the application, the positive specimen with a mixing ratio of 1:2 showed the darkest line, subsequently those with mixing ratios of 1:4 and 1:1 showed the darker lines to equal extent, followed by those with mixing ratios of 1:0.5 and 1:8 in this order. In contrast, specimens containing the anti-TP antibody negative serum with individual mixing ratios showed no line.

The result shows that the detection sensitivity can easily be changed by changing the mixing ratio of the oligonucleotide-labeled avidin to the biotin-labeled antigen.

EXAMPLE 14

Preparation of Oligonucleotide-labeled IgG for Detection of HBs Antigen

The oligonucleotide Pair 1- was introduced into a rabbit IgG in a similar manner to that in Example 2.

EXAMPLE 15

Preparation of Oligonucleotide-labeled Mouse Anti-HBs-Fab' for the Detection of HBs Antigen

To a 0.1 M citrate buffer pH 3.5 containing 10 mg of a monoclonal anti-HBs-IgG, pepsin (available from Boehringer Mannheim) was added to 5% of the IgG and was allowed to react at 37 ° C for a predetermined time. After the reaction, the resulting mixture was applied to a Ultrogel AcA44 column (trade name; available from Biosepra) equilibrated with 0.1 M phosphate buffer pH 6.0 to thereby collect an F(ab')₂ fraction.

A total of 5.2 mg of the above-obtained F(ab')₂ was concentrated by ultrafiltration using YM30 (available from Millipore Corporation), the concentrate was reduced at 37° C for 90 minutes with mercaptoethylamine (available from Nakarai Tesque, Inc.) in a final concentration of 0.1 M. The resulting mixture was then applied to a Sephadex G-25 column (trade name, available from Amersham Pharmacia Biotech Company (Amersham Biosciences)) and thereby yielded 2.6 mg of a mouse anti-HBs-Fab'.

Separately, to 0.1 M MOPS buffer pH 7.0 containing 191 nmol of Pair 1+ dissolved therein and 5 mM EDTA, a 100-fold by mole excess of EMCS with respect to the oligonucleotide was added and was allowed to react at 37° C for 30 minutes. After the reaction, the resulting mixture was subjected to precipitation with ethanol and washing according to a conventional procedure and thereby yielded a maleimide-group-introduced oligonucleotide.

The maleimide-group-introduced oligonucleotide was allowed

to react with the mouse anti-HBs-Fab' at 37° C for 1 hour, was then applied to a Ultrogel AcA44 column and thereby yielded "Pair 1+"-labeled mouse anti-HBs-Fab'.

EXAMPLE 16

5 Preparation of Nitrocellulose Membrane for the Simultaneous Detection of HBs Antigen and Anti-TP Antibody

The "Pair 8+"-labeled rabbit IgG prepared in Example 2 and the "Pair 1+"-labeled rabbit IgG prepared in Example 14 were diluted to a concentration of 1 mg/ml and were charged into
10 XY3000 (trade name, available from BioDot, Inc.), respectively. The "Pair 8+"-labeled rabbit IgG and the "Pair 1+"-labeled rabbit IgG were applied to a nitrocellulose membrane lined with PET at positions of 35 mm and 30 mm from a starting point, respectively. After drying, the applied membrane was blocked
15 with Blockace (trade name, available from Snow Brand Milk Products Co., Ltd.) at room temperature for 3 hours, was washed with distilled water and was then dried. A cellulose filter paper (available from Whatman Inc. under the trade name of WF1.5 or from Advanced Microdevices Pvt. Ltd. (mdi)) was attached to
20 one end of the dried article, and a conjugate release pad (available from Advanced Microdevices Pvt. Ltd. (mdi)) to the other end. In this procedure, part of them were overlapped with each other. The resulting article was cut to 5 mm in width and was housed in a housing.

25 EXAMPLE 17

Analysis on Possibility of Simultaneous Detection of HBs Antigen and Anti-TP Antibody

The possibility of simultaneous detection of a HBs antigen and an anti-TP antibody was analyzed in the following manner.

Specifically, the oligonucleotide-labeled TP 17K antigen with a mixing ratio of 1:4 prepared in Example 12 was diluted ten folds, and 1 ul of the diluted solution was added to 5 ul of the colloidal gold-labeled TP antigen prepared in Example 9, 5 ul of a biotin 1 mg/ml solution and 90 ul of an anti-TP antibody negative and HBs antigen negative serum. To the resulting mixture, 90 ng of the oligonucleotide-labeled mouse anti-HBs-Fab' prepared in Example 15 and 9 ul of a colloidal gold-labeled anti-HBs antibody (available from British Biocell International) were added. Four bottles of this reaction system were prepared. A total of 5 ul of an anti-TP antibody positive serum was added to one bottle, 5 ul of an anti-TP antibody negative and HBs antibody negative serum was added to another, and 5 ul of sera containing 50 ng/ml and 220 ug/ml of HBs antigen were added to the remained two bottles, respectively, and the resulting mixtures were immediately mixed, and the total amounts of the resulting mixtures were applied to the sample supply opening of the housing containing the "Pair 8-" -labeled rabbit IgG-binding and "Pair 1-" -labeled rabbit IgG-binding nitrocellulose membrane prepare in Example 16, respectively.

Twenty minutes after the application, the sample containing the anti-TP antibody negative and HBs antigen negative serum showed no line. The sample containing the anti-TP antibody positive serum showed a line only in a portion where the Pair

8- oligonucleotide was applied, and the samples each containing the HBs antigen showed a line only in a portion where the Pair 1- oligonucleotide was applied. Each of them was independently detected on the same nitrocellulose membrane.

5 EXAMPLE 18

Preparation of Biotin-labeled TP 17K Antigen

To 0.5 ml of 0.1 M phosphate buffer pH 7.0 containing 500 ug of TP 17K antigen (available from Lee Laboratories) and 5 mM EDTA, a 100-fold by mole excess of an NHS-biotin was added with respect to the TP 17K antigen and was allowed to react at 37° C for 1 hour. After the reaction, the resulting mixture was applied to a Sephadex G-25 column and thereby yielded a biotin-labeled TP 17K antigen with a recovery of the TP 17K antigen of 75%. The ultimate recovery of the biotin-labeled TP 17K antigen was 75%, since it was prepared merely by mixing the TP 17K antigen with the oligonucleotide-labeled avidin prepared in Example 10, as described in Example 12.

COMPARATIVE EXAMPLE

20 Preparation of Oligonucleotide-labeled TP 17K Antigen Through Direct Labeling of TP 17K Antigen

To 0.1 M phosphate buffer pH 7.4 containing 473 μ g of a TP 17K antigen (available from Lee Laboratories) and 5 mM EDTA, a 20-fold by mole excess of SATA was added with respect to the TP 17K antigen and was allowed to react at 37 ° C for 90 minutes. Hydroxylamine and Tris buffer were added to and were allowed to react with the resulting mixture. The resulting reaction mixture was then applied to a Sephadex G-25 column (trade name,

available from Amersham Pharmacia Biotech Company (Amersham Biosciences)) and thereby yielded 200 μ g of an SH-group-introduced TP 17K antigen (recovery: 42.2%). Separately, to 0.1 M MOPS buffer pH 7.85 containing 133 nmol of Pair 7+ and 5 mM EDTA, a 200-fold by mole excess of EMCS with respect to the oligonucleotide Pair 7+ was added and was allowed to react at 37 ° C for 2 hours. The resulting oligonucleotide was subjected to precipitation with ethanol and rinsing according to a conventional procedure and thereby yielded 105 nmol of a maleimide-group-introduced oligonucleotide.

A total of 200 mg of the SH-group-introduced TP 17K antigen and 105 nmol of the maleimide-group-introduced oligonucleotide were mixed and were allowed to react with each other at 37° C for 2 hours. The resulting mixture was then applied to an Ultrogel AcA44 column and thereby yielded 150 μ g of an oligonucleotide-labeled TP 17K antigen (recovery: 32%).

In comparison between Example 18 and Comparative Example, the product according to Comparative Example showed a recovery of 32%, and the product according to Example 18 showed a recovery of 75%, indicating that the recovery was significantly improved in Example 18.

EXAMPLE 19

Preparation of HCV Core 2, Core 3, Core 6 and Core 10-Binding colloidal gold-labeled Streptavidin

A total of 500 μ l of a colloidal gold-labeled streptavidin (available from British Biocell International) prepared to show an absorbance at 530 nm of 10 was mixed with 2 μ g of a

biotinylated HCV core 2, 2 μ g of a biotinylated HCV core 3, 2 μ g of a biotinylated HCV core 6, and 2 μ g of a biotinylated HCV core 10 (each available from Innogenetics NV) and was allowed to react at 37° C for 1 hour. After the reaction, a HCV core 2, core 3, core 6 and core 10-binding colloidal gold-labeled streptavidin was recovered as a precipitate by centrifugation. The precipitate was suspended in 20 mM sodium borate buffer pH 8.0 containing 2% bovine serum albumin and was stored at 4° C.

10 EXAMPLE 20

Oligonucleotide-labeled HCV Core 2, Core 3, Core 6, or Core 10-binding Streptavidin

To 100 μ g of the "Pair 8+"-labeled streptavidin prepared in Example 4, 30 μ g each of HCV core 2, core 3, core 6, or core 10 was added, respectively, was allowed to react at 37 ° C for 1 hour, the resulting mixtures were subjected to ultrafiltration and thereby yielded a "Pair 8+"-labeled HCV core 2-binding streptavidin, a "Pair 8+"-labeled HCV core 3-binding streptavidin, a "Pair 8+"-labeled HCV core 6-binding streptavidin, and a "Pair 8+"-labeled HCV core 10-binding streptavidin, respectively.

EXAMPLE 21

Preparation of Oligonucleotide-labeled HCV Core 2, Core 3, Core 6 and Core 10-binding Streptavidin

To 100 μ g of the "Pair 8+"-labeled streptavidin prepared in Example 4, 30 μ g each of HCV core 2, core 3, core 6 and core 10 were added concurrently and were allowed to react at

37 ° C for 1 hour. The resulting mixture was subjected to ultrafiltration and thereby yielded a "Pair 8+"-labeled HCV core 2, core 3, core 6 and core 10-binding streptavidin.

EXAMPLE 22

5 Reactivity with Anti-HCV antibody Positive Serum

10 The reactivity of the above-prepared articles was analyzed in the same manner as in Example 8. Specifically, a combination (Combination A) was prepared by mixing the HCV core 2, core 3, core 6 and core 10-binding colloidal gold-labeled streptavidin prepared in Example 19 with 37.5 ng of the Pair 8+"-labeled HCV core 2-binding streptavidin, 37.5 ng of the Pair 8+"-labeled HCV core 3-binding streptavidin, 37.5 ng of the Pair 8+"-labeled HCV core 6-binding streptavidin, and 37.5 ng of the Pair 8+"-labeled HCV core 10-binding streptavidin each prepared in Example 20. Another combination (Combination B) was prepared by mixing 5 ul of the HCV core 2, core 3, core 6 and core 10-binding colloidal gold-labeled streptavidin prepared in Example 19 with 150 ng of the "Pair 8+"-labeled HCV core 2, core 3, core 6 and core 10-binding streptavidin prepared in Example 21. The reaction with an anti-HCV antibody positive serum was observed 5 minutes and 7 minutes into the test in Combination A and in Combination B, respectively. In this connection, the reaction was observed 15 minutes into the test in Example 8. These results show that a reaction system including plural types of peptides bound to a labeled streptavidin exhibits higher reactivity than a reaction system including a single type of peptide bound to a labeled

streptavidin.

INDUSTRIAL APPLICABILITY

5 The kits and methods for detection or quantification of the present invention can control the sensitivity of reagents.

For example, the kits of the present invention can control the amount of the ligand L1 or the ligand L2 per molecule of the receptor I and receptor II. This configuration can control the sensitivity of measurement. Specifically, a substance
10 having plural binding points to the substance B1 can be used as the substance R1, and thereby plural pieces of the ligand L1 can be bound to the substance R1 through the medium of the substance B1. Likewise, a substance having plural binding points to the substance B2 can be used as the substance R2, and
15 thereby plural pieces of the ligand L2 or the ligand L3 can be bound to the substance R2 through the medium of the substance B2.

Specifically, when avidin or streptavidin is used as the substance R1 or R2, the avidin or streptavidin has four binding
20 points per its molecule with respect to biotin as the substance B1, and the ratio of the introduced ligand L1 can be changed from one to four per molecule of avidin or streptavidin. The control of the inducing ratio can ensure the control of the reactivity and sensitivity. In addition, the sensitivity can be
25 controlled by the compound of L-B1-R1-M and the compound of L-B1-R2-B3, ensuring the control of the sensitivity to further detail.

By selecting combinations of substances having different binding capabilities as the bond element B3 and the anti-bond element R3, the intensity of binding capability can be changed, and the sensitivity of reagents can be controlled.

5 When the bond element B3 and the anti-bond element R3 are oligonucleotides of RNA and/or DNA at least having partly complementary sequences and are bound to each other by action of their complementarity, the intensity of binding capability can be controlled, and thereby the sensitivity of the reagent can be
10 controlled by changing the lengths and/or homology of the oligonucleotides.

In the kits of the present invention, the plural ligands to be bound to the substance R1 or R2 may have the same reactivity or may have plural types of reactivity. In the latter case, the
15 resulting kit has higher reactivity to the analyte A than a kit including ligands having the same reactivity.

According to the kits and methods for detection or measurement of the present invention, when plural oligonucleotides of, for example, RNAs or DNAs are used as the
20 bond element B3 and the anti-bond element R3 to be solid-phased, and the resulting complex is bound to the solid phase through at least partly complementary binding, oligonucleotides having different sequences can be used as the oligonucleotides. The invented kits and methods for detection or measurement can
25 therefore simultaneously detect or quantify plural analytes corresponding to the plural types of the oligonucleotides.

What Is Claimed Is:

1. A kit for detecting or measuring an analyte having bivalent or higher binding capability, the kit comprising the following receptor I, receptor II, and solid phase conjugate for capturing a complex comprising the receptor I, the receptor II and the analyte:

1) a receptor I represented by a compound L1-B1-R1-M comprising a compound R1-M bound to a compound L1-B1, the compound R1-M comprising a substance R1 bound to a marker M, the substance R1 being capable of binding to a substance B1, the compound L1-B1 comprising a ligand L1 bound to the substance B1, the ligand L1 being capable of binding to the analyte A;

2) a receptor II represented by a compound L2-B2-R2-B3 comprising a compound L2-B2 previously bound to a compound R2-B3, the compound L2-B2 comprising a ligand L2 and a substance B2 introduced into the ligand L2, the ligand L2 being capable of binding to the analyte A, the substance B2 having different binding capability from the analyte A, the compound R2-B3 comprising a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3 having binding capability different from the substance B2; and

3) solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3.

2. A kit for detecting or measuring an analyte A having bivalent or higher binding capability to a ligand L3 constituting part of a protein P, the kit comprising the following receptor I, receptor II, and solid phase conjugate for capturing a complex comprising the receptor I, the receptor II and the analyte:

1) a receptor I represented by a compound P-M comprising the protein P bound to a marker M;

2) a receptor II represented by a compound P-B2-R2-B3 or L3-B2-R2-B3 comprising a compound P-B2 or L3-B2 bound to a compound R2-B3, the compound P-B2 or L3-B2 comprising the protein P or the ligand L3 and a substance B2 introduced into the protein P or the ligand L3, the substance B2 having different binding capability from the analyte A, the compound R2-B3 comprising a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3 having binding capability different from the substance B2; and

3) solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3.

3. A kit for detecting or measuring an analyte A having bivalent or higher binding capability, the kit comprising the following receptor I, receptor II, and solid phase conjugate for capturing a complex comprising the receptor I, receptor II

and analyte:

1) a receptor I represented by a compound L1-B1-R1-M comprising a compound R1-M bound to a compound L1-B1, the compound R1-M comprising a substance R1 bound to a marker M, the substance R1 being capable of binding to a substance B1, the compound L1-B1 comprising a ligand L1 bound to the substance B1, the ligand L1 being capable of binding to the analyte A;

2) a receptor II represented by a compound L2-B3, the compound L2-B3 comprising a ligand L2 previously bound to a bond element B3, the ligand L2 being capable of binding to the analyte A, the bond element B3 having different binding capability from analyte A; and

3) solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3.

4. The kit according to claim 1 or 3, wherein the substance R1 has a plurality of binding points with respect to the substance B1, and a plurality of the ligands L1 are bound to the substance R1 by the medium of the substance B1.

5. The kit according to claim 4, wherein each of the plurality of the ligands L1 has plural types of reactivity.

6. The kit according to claim 1 or 2, wherein the substance R2 has a plurality of binding points with respect to the

substance B2, and a plurality of the ligands L2 or ligands L3 are bound to the substance R2 by the medium of the substance B2.

7. The kit according to claim 6, wherein each of the
5 plurality of the ligands L2 or ligands L3 has plural types of reactivity.

8. The kit according to any one of claims 1 to 3, wherein
10 the solid phase conjugate is at least independent of the receptor II.

9. The kit according to any one of claims 1 to 3, wherein
the analyte A having bivalent or higher binding capability is a
substance selected from the group consisting of DNAs, RNAs,
15 antigens, and antibodies.

10. The kit according to any one of claims 1 to 3, wherein
the ligand L1, ligand L2 or ligand L3 is a substance selected
from the group consisting of DNA, RNA, antigen, antibody,
20 lectin, glycoprotein, and sugar.

11. The kit according to claim 1 or 3, wherein the ligand
L1 and the ligand L2 are the same substance.

12. The kit according to any one of claims 1 to 3, wherein
25 the ligands L1 and L2 and the ligand L3 are substances having different sequences with one another.

13. The kit according to any one of claims 1 to 3, wherein the binding capability between the substance B1 and the substance R1 or between the substance B2 and the substance R2 is represented by a dissociation constant of from 10^{-8} to 10^{-16} (M).

14. The kit according to any one of claims 1 to 3, wherein the substance B1 and/or the substance B2 is biotin.

15. The kit according to any one of claims 1 to 3, wherein the substance B1 and/or the substance B2 is a substance selected from the group consisting of DNA, RNA, antigen, antibody, lectin, glycoprotein, and sugar.

16. The kit according to any one of claims 1 to 3, wherein the substance R1 and/or the substance R2 is a substance selected from the group consisting of streptavidin and avidin.

17. The kit according to any one of claims 1 to 3, wherein the substance R1 and/or the substance R2 is a substance selected from the group consisting of antigen, antibody, DNA, RNA, lectin, glycoprotein, and sugar.

18. The kit according to claim 1, wherein the substance R1 and the substance R2 are the same substance.

19. The kit according to claim 1, wherein the substance R1 and the substance R2 are different substances.

20. The kit according to any one of claims 1 to 3, wherein
5 the bond element B3 is bondable to the anti-bond element R3 by complementary binding of at least part of DNAs or RNAs.

21. The kit according to any one of claims 1 to 3, wherein
10 the marker M is a substance selected from the group consisting of coloring dye, fluorescent dye, luminescent substance, metal colloid, latex, liposome, radioactive isotope, enzyme, DNA, and RNA.

22. The kit according to any one of claims 1 to 3, wherein
15 the solid phase is a substance selected from the group consisting of polystyrene, nitrocellulose, nylon, cellulose, and glass.

23. A method for detecting or measuring an analyte A having
20 bivalent or higher binding capability, the method comprising the steps of:

bringing 1) the analyte A into contact with:

2) a receptor I represented by a compound L1-B1-R1-M
comprising a compound R1-M bound to a compound L1-B1, the
25 compound R1-M comprising a substance R1 bound to a marker M,
the substance R1 being capable of binding to a substance B1, the
compound L1-B1 comprising a ligand L1 bound to the substance

B1, the ligand L1 being capable of binding to the analyte A; and

3) a receptor I1 represented by a compound L2-B2-R2-B3 comprising a compound L2-B2 previously bound to a compound R2-B3, the compound L2-B2 comprising a ligand L2 and a substance B2 introduced into the ligand L2, the ligand L2 being capable of binding to the analyte A, the substance B2 having different binding capability from the analyte A, the compound R2-B3 comprising a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3 having binding capability different from the substance B2, to react with one another to thereby form a complex;

4) allowing a solid phase conjugate to capture the formed complex, the solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3, and

5) detecting or measuring the marker M in the captured complex.

24. A method for detecting or measuring an analyte A having bivalent or higher binding capability to a ligand L3 constituting part of a protein P, the method comprising the steps of:

bringing 1) the analyte A into contact with:

2) a receptor I represented by a compound P-M comprising the protein P bound to a marker M; and

3) a receptor II represented by a compound P-B2-R2-B3 or L3-B2-R2-B3 comprising a compound P-B2 or L3-B2 bound to a compound R2-B3, the compound P-B2 or L3-B2 comprising the protein P or the ligand L3 and a substance B2 introduced into the protein P or the ligand L3, the substance B2 having different binding capability from the analyte A, the compound R2-B3 comprising a substance R2 bound to a bond element B3, the substance R2 being capable of binding to the substance B2, the bond element B3 having binding capability different from the substance B2, to react with one another to thereby form a complex,

4) allowing a solid phase conjugate to capture the formed complex, the solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3, and

5) detecting or measuring the marker M in the captured complex.

25. A method for detecting or measuring an analyte A having bivalent or higher binding capability, the method comprising the steps of:

bringing 1) the analyte A into contact with:

2) a receptor I represented by a compound L1-B1-R1-M comprising a compound R1-M bound to a compound L1-B1, the compound R1-M comprising a substance R1 bound to a marker M, the substance R1 being capable of binding to a substance B1, the compound L1-B1 comprising a ligand L1 bound to the substance

B1, the ligand L1 being capable of binding to the analyte A; and

3) a receptor I1 represented by a compound L2-B3, the compound L2-B3 comprising a ligand L2 previously bound to a bond element B3, the ligand L2 being capable of binding to the analyte A, the bond element B3 having different binding capability from analyte A, to react with one another to thereby form a complex,


4) allowing a solid phase conjugate to capture the formed complex, the solid phase conjugate represented by a compound R3-solid phase, the compound R3-solid phase comprising a anti-bond element R3 bound to a solid phase, the anti-bond element R3 being capable of binding to the bond element B3, and

5) detecting or measuring the marker M in the captured complex.


ABSTRACT

A kit and a production method therefor are provided, which kit can be produced in a high yield, can easily label ligands and can control the sensitivity of reagents. A kit for detecting or measuring an analyte having bivalent or higher binding capability includes the following receptor I, receptor II, and solid phase conjugate. Receptor I: L1-B1-R1-M including a compound R1-M bound to a compound L1-B1, in which the compound R1-M includes a substance R1 being capable of binding to a substance B1 bound to a marker M, the compound L1-B1 includes a ligand L1 and the substance B1. Receptor II: L2-B2-R2-B3 including a compound L2-B2 previously bound to a compound R2-B3, in which the compound L2-B2 includes a ligand L2 capable of binding to the analyte A and a substance B2 being introduced into the ligand L2 and having different binding capability from the analyte A, the compound R2-B3 includes a substance R2 capable of binding to the substance B2 and being bound to a bond element B3 having different binding. Solid phase conjugate: R3-solid phase including a anti-bond element R3 bound to a solid phase, in which the anti-bond element R3 is capable of binding to the bond element B3.


F i g . 1



F i g . 2



F i g . 3



F i g . 4




Fig. 5




Fig. 6




Fig. 7





Fig. 8



COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

This declaration is of the following type:

- ☐ original
- ☐ design
- ☐ supplemental
- ☒ national stage of PCT
- ☐ divisional
- ☐ continuation
- ☐ continuation-in-part (CIP)

My residence, post office address and citizenship are as stated next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed for and for which a patent is sought on the invention entitled:

KITS AND METHODS FOR DETECTING OR MEASURING ANALYTES

the specification of which

- ☐ is attached hereto
- ☐ was filed on _____, as
Application Serial No. _____
and was amended on _____
(if applicable)
- ☒ was described and claimed in PCT International application
No. PCT/JP 00/06187 filed on 11, September 2000
and as amended under PCT Article 19 on _____
(if any).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any Amendment referred to above.

I acknowledge duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Sec. 1.56.

- ☐ In compliance with this duty there is attached an information disclosure statement. 37 CFR 1.97.

I hereby claim foreign priority benefits under Title 35, United States Code, Sec. 119, of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent of inventor's certificate having a filing date before that of the application on which priority is claimed:

[] no such applications have been filed
[X] such applications have been filed as follows.

Prior Foreign Application(s)

<u>11-258771</u>	<u>Japan</u>	<u>13 / September / 1999</u>	[X]	[]
(Number)	(Country)	(day/month/year filed)	Yes	No
<u> </u>	<u> </u>	<u> </u>	[]	[]
(Number)	(Country)	(day/month/year filed)	Yes	No

I hereby claim the benefit under Title 35, United States Code, Sec. 120 of any United States application(s) listed below, and insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, Sec. 112, I acknowledge the duty to disclose all information known to be material to patentability as defined in Title 37, Code of Federal Regulations, Sec. 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application:

<u> </u>	<u> </u>	<u> </u>
(Application Serial No.)	(Filing Date)	(patented, pending, abandoned)
<u> </u>	<u> </u>	<u> </u>
(Application Serial No.)	(Filing Date)	(patented, pending, abandoned)

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agents to prosecute this application and transact all business in the Patent and Trademark Office connected therewith.

George A. Loud, Reg. No. 25,814
Anthony M. Lorusso, Reg. No. 25,059
Donald W. Marks, Reg. No. 24,218

send correspondence to

Direct telephone calls
TO:

LORUSSO & LOUD
3137 Mount Vernon Avenue
Alexandria, VA 22305

(703) 739-9393

I hereby declare all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor 1-0 Yuichi Oku

Inventor's signature Yuichi Oku Feb. 22, 2002 (Date)

Residence Ibaraki 307-0036 Japan SPX

Citizenship Japanese

Post Office Address C/O NISSUI PHARMACEUTICAL CO., LTD.,
RESEARCH DEPARTMENT, 1075-2 Hokunanmoro,
Yuuki-shi, Ibaraki 307-0036 Japan

Full name of second joint inventor, if any 2-0 Hisanori Kamiya

Inventor's signature Hisanori Kamiya Feb. 22, 2002 (Date)

Residence Ibaraki 307-0036 Japan SPX

Citizenship Japanese

Post Office Address C/O NISSUI PHARMACEUTICAL CO., LTD.,
RESEARCH DEPARTMENT, 1075-2 Hokunanmoro,
Yuuki-shi, Ibaraki 307-0036 Japan

Full name of third joint inventor, if any 3-0 Kumiko Shinohara

Inventor's signature Kumiko Shinohara Feb. 22, 2002 (Date)

Residence Ibaraki 307-0036 Japan SPX

Citizenship Japanese

Post Office Address C/O NISSUI PHARMACEUTICAL CO., LTD.,
RESEARCH DEPARTMENT, 1075-2 Hokunanmoro,
Yuuki-shi, Ibaraki 307-0036 Japan

Full name of fourth joint inventor, if any 4-D Yusuke Sibahara

Inventor's signature Yusuke Sibahara Feb. 21, 2002 (Date)

Residence Ibaraki 307-0036 Japan SPX

Citizenship Japanese

Post Office Address C/O NISSUI PHARMACEUTICAL CO., LTD.,
RESEARCH DEPARTMENT, 1075-2 Hokunanmoro,
Yuuki-shi, Ibaraki 307-0036 Japan

Full name of fifth joint inventor, if any 5-D Yoshihiko Uesaka

Inventor's signature Yoshihiko Uesaka Feb. 22, 2002 (Date)

Residence Ibaraki 307-0036 Japan SPX

Citizenship Japanese

Post Office Address C/O NISSUI PHARMACEUTICAL CO., LTD.,
RESEARCH DEPARTMENT, 1075-2 Hokunanmoro,
Yuuki-shi, Ibaraki 307-0036 Japan

204060 T9907001

SEQUENCE LISTING

<110> Nissui Pharmaceutical Co., Ltd.

<120> Kit for detection or measurement of material to be
measured and method for detection or measurement

<130> NSM0037

<160> 5

<210> 1

<211> 20

<212> DNA

<223> The oligonucleotide with the amino group in the 5'
terminal.

<400> 1

gaattcccg g gatccgtcg 20

<210> 2

<211> 20

<212> DNA

<223> The oligonucleotide with the amino group in the 5'
terminal.

<400> 2

cgacggatcc ccgggaattc 20

<210> 3

<211> 20

<212> DNA

<223> The oligonucleotide with the amino group in the 5'
terminal.

<400> 3

aacggaatct aatcaggagg 20

<210> 4

<211> 20

<212> DNA

<223> The oligonucleotide with the amino group in the 5' terminal.

<400> 4

cctcctgatt agattccgtt 20

<210> 5

<211> 20

<212> DNA

<223> The oligonucleotide with the amino group in the 5' terminal.

<400> 5

ccgactacag aagaggagaa 20